

Decoding Calcium Signaling Across the Nucleus

Calcium (Ca^{2+}) is an important multifaceted second messenger that regulates a wide range of cellular events. A Ca^{2+} -signaling toolkit has been shown to exist in the nucleus and to be capable of generating and modulating nucleoplasmic Ca^{2+} transients. Within the nucleus, Ca^{2+} controls cellular events that are different from those modulated by cytosolic Ca^{2+} . This review focuses on nuclear Ca^{2+} signals and their role in regulating physiological and pathological processes.

André G. Oliveira,¹
Erika S. Guimarães,^{1,2}
Lídia M. Andrade,¹
Gustavo B. Menezes,³ and
M. Fatima Leite¹

¹Department of Physiology and Biophysics, Federal University of Minas Gerais, Belo Horizonte, Brazil; ²Molecular Medicine, School of Medicine, Federal University of Minas Gerais, Belo Horizonte, Brazil; and ³Department of Morphology, Federal University of Minas Gerais, Belo Horizonte, Brazil
leitemd@ufmg.br

The role of calcium (Ca^{2+}) signaling in regulating a variety of cellular processes has been widely described, showing the multifunctional nature of this ion. Even though the mechanisms that allow Ca^{2+} to modulate a wide range of events have not been completely elucidated, it is appreciated that the amplitude, the spatiotemporal profile (e.g., single transient, wave, oscillation), as well as the intracellular compartments where Ca^{2+} signals occur are all involved in orchestrating Ca^{2+} increases that correspond to specific cellular functions mediated by Ca^{2+} . In this review, we will briefly discuss the machinery that regulates intracellular Ca^{2+} signaling and then highlight mechanisms of Ca^{2+} signaling in the nucleus as well as cellular events that are modulated by nucleoplasmic Ca^{2+} .

Intracellular Ca^{2+} -Signaling Toolkit

Although the variety of biological processes regulated by Ca^{2+} can be partially explained by the spatiotemporal organization of Ca^{2+} increases within the cell, a crucial question in this signaling cascade is how cells can generate, regulate, and coordinate such Ca^{2+} -evoked responses. The remarkable plasticity of Ca^{2+} signaling emerges not only from the origin of the intracellular Ca^{2+} signals (i.e., endoplasmic reticulum, mitochondria, or nucleus) but also from the molecular toolkit expressed by different cell types, as well as from the interplay between the cellular compartments. Below, we will briefly review some of the components of the Ca^{2+} toolkit that have been described to date. Although we will not discuss the components at length and we will not present a complete list of the Ca^{2+} signaling molecules, the discussion will provide a sense of the variety of components of the Ca^{2+} signaling network that can be used by different cell types to generate a specific cellular event.

Intracellular Ca^{2+} signals can be initiated by the influx of Ca^{2+} from the extracellular environment or by the mobilization of Ca^{2+} from intracellular

stores. The latter depends mainly on the binding of hormones or growth factors to specific receptor recruiting and generating a variety of second messengers that, in turn, will control Ca^{2+} mobilization from different intracellular stores (reviewed in Refs. 10, 87). One of the major signaling cascades triggered by the activation of transmembrane receptors involves phospholipase C (PLC), which generates inositol-1,4,5-trisphosphate (InsP_3). InsP_3 diffuses throughout the cell to bind to its specific receptor (InsP_3R), which is a ligand-gated Ca^{2+} release channel (reviewed in Refs. 19, 30). Three isoforms of InsP_3R have been described, and although these receptors are mainly located in the endoplasmic reticulum, they can also be detected at the plasma membrane (25), portions of the Golgi complex (83), acidic stores (35), and within the nucleus (29, 45, 59). Moreover, the expression pattern of InsP_3R isoforms varies among different cell types (108). Although InsP_3 is required to activate InsP_3R , the opening of the channel can be modulated by Ca^{2+} itself, ATP, and proteins such as Chromogranin-B (reviewed in Refs. 19, 24, 30). It is also known that the affinity of InsP_3R for InsP_3 is highest for $\text{InsP}_3\text{R-II}$, intermediate in $\text{InsP}_3\text{R-I}$, and lowest for $\text{InsP}_3\text{R-III}$ (52, 74).

The intracellular increase in Ca^{2+} concentrations itself may activate a second class of receptors: the ryanodine receptors (RyR). Similar to InsP_3R , RyR forms channels that are permeable to Ca^{2+} . So far, three isoforms of RyR have been identified (reviewed in Refs. 57, 112). RyR-I is abundantly expressed in skeletal muscle but also is found in cardiac muscle, cerebellum, and other tissues (33, 40, 75). RyR-II is the major isoform found in cardiac muscle (71, 79) but is also expressed in other tissues (33, 55, 72, 96), whereas RyR-III is expressed within hippocampal neuron cerebellum as well as in smooth muscle cells of many organs (33, 40, 46). Ca^{2+} is the major trigger of RyR opening, but some RyR channel subtypes are also activated by cyclic ADP-ribose (cADPR) (70, 100). Moreover, FK506 binding protein,

magnesium, calmodulin, and a number of small molecules such as ATP are able to modulate the activity of RyRs (Ref. 68; reviewed in Refs. 11, 112).

In addition to the well known mechanisms above, a growing body of evidence shows that members of a third and a fourth class of intracellular Ca^{2+} channels can be involved in controlling Ca^{2+} release from organelles within the cell. One family of these receptors, named two pore channels (TPCs), acts as Ca^{2+} channels and is responsible for the mobilization of Ca^{2+} from acidic organelles in response to stimulation by nicotinic acid adenine dinucleotide phosphate (NAADP) (15, 34). There are three isoforms of TPCs in mammals, with TPC1 and TPC3 expressed in endosomes and TPC2 present primarily in lysosomes (15, 34, 92). It has been shown that TPC-induced Ca^{2+} release may trigger further Ca^{2+} release through the activation of juxtaposed InsP_3R or RyR, suggesting a broader role of TPCs in the regulation of cellular activities (Refs. 16, 34; reviewed in Ref. 102).

Another channel shown to be involved in intracellular Ca^{2+} signaling is the TRP2 that belongs to the transient receptor potential (TRP) superfamily of ion channels (Refs. 53, 54; reviewed in Refs. 5, 104). Besides its location at the plasma membrane and the primary cilium, TRP2 is also found in high concentrations in the endoplasmic reticulum membrane functioning as an intracellular Ca^{2+} -release channel that augments InsP_3 -induced intracellular Ca^{2+} release (54).

Within a cell, Ca^{2+} signaling may be seen as a shifting balance that controls cell physiology. When this signaling pathway is activated but tightly controlled, Ca^{2+} regulates a broad spectrum of biological processes, including secretion, motility, gene transcription, proliferation, differentiation, contraction, cell death, and others (10). Therefore, it is important that the cells express a toolkit not only to allow Ca^{2+} release from intracellular stores but also to reestablish the intracellular free Ca^{2+} to baseline levels. To accomplish this, cytosolic Ca^{2+} can be 1) extruded into the extracellular environment by the action of the plasma membrane Ca^{2+} -ATPase (PMCA) or the sodium Ca^{2+} exchanger (NCX), or 2) pumped back into the lumen of the Ca^{2+} stores, such as endoplasmic reticulum and mitochondria, by the action of proteins like the sarco-/endoplasmic Ca^{2+} -ATPase (SERCA) and mitochondrial Ca^{2+} uptake 1 (MCU1) (Ref. 82; reviewed in Refs. 12, 84).

Although understanding the components of the intracellular Ca^{2+} toolkit constitutes a powerful strategy to familiarize the reader with the diversity and peculiarities of the molecular repertoire involved in intracellular Ca^{2+} signaling, the reader should be reminded that these components act in a coordinated fashion to regulate specific cellular

functions triggered by increases in Ca^{2+} . Also, despite the broad expression of the components of the Ca^{2+} toolkit, Ca^{2+} signaling may occur in distinct microdomains of the cell (9). Evidence showing components of the Ca^{2+} toolkit in the nucleus identifies this cellular compartment as a pivotal microdomain for independent and localized Ca^{2+} signaling. In the following sections, we will describe the mechanisms underlying nuclear Ca^{2+} signaling as well as its relevance for physiological and pathological processes.

Nucleoplasmic Ca^{2+} Signaling

There are several reports in the literature suggesting that nuclear Ca^{2+} signals can result from passive diffusion of Ca^{2+} from the cytosol to the nucleoplasm (14, 48, 61, 98). This view evolved from observations made by numerous groups showing not only the existence of nuclear-cytosolic Ca^{2+} gradients (94, 107) but also the presence of several components of the Ca^{2+} toolkit within the nucleus capable of autonomously controlling nuclear Ca^{2+} homeostasis (reviewed in Refs. 37, 87, 91).

For instance, InsP_3 -sensitive pools (77), as well as a high-affinity, InsP_3 binding site, were demonstrated to exist in isolated liver nuclei, with InsP_3 being able to release $^{45}\text{Ca}^{2+}$ directly from isolated nuclei (63). Also using isolated rat liver nuclei, it was shown that ATP increases the free Ca^{2+} concentration in the nucleoplasm (76). However, it was not initially clear which intranuclear structures contained Ca^{2+} nor into which intranuclear compartments Ca^{2+} was released by InsP_3 . Indeed, the first report demonstrating that InsP_3 is able to cause the release of Ca^{2+} from the nuclear envelope into the nucleoplasm came a few years later. It was then shown that the nuclear envelope actively takes up Ca^{2+} via a Ca^{2+} -activated ATPase and that InsP_3 releases Ca^{2+} from this envelope store into the nucleoplasm (39). A subsequent study, in which InsP_3 was injected in the nucleus of *Xenopus laevis* oocytes, also demonstrated that InsP_3 causes an increase in nuclear Ca^{2+} , even when cytosolic InsP_3Rs were blocked (48). Moreover, photorelease of caged InsP_3 in the nucleus of starfish oocytes was shown to trigger nucleoplasmic Ca^{2+} increase (94). Like InsP_3 , cADPR also could raise Ca^{2+} in isolated hepatocyte nuclei (1, 39) and induce Ca^{2+} oscillations in the nucleus of starfish oocytes (94). These findings illustrate some of the initial evidence that supported the hypothesis that the nucleus could autonomously trigger nucleoplasmic Ca^{2+} transients. This hypothesis was strengthened by additional data showing that the inner and/or the outer membranes of the nuclear envelope express several components of the Ca^{2+} toolkit. In particular, it was shown that the nuclear envelope

accumulates Ca^{2+} via a Ca^{2+} -ATPase pump (SERCA) (56, 76) located in its outer leaflets (38, 49) and a $\text{Na}^+/\text{Ca}^{2+}$ -exchanger (109, 110) located in its inner membrane (109), and releases Ca^{2+} via channels that are sensitive to InsP_3 (39, 76), cADPR (1, 39), and NAADP (36).

There are reports that the InsP_3R (17, 49, 101) and the RyR (93) can be present on both leaflets of the nuclear envelope, whereas ADP-ribosyl (CD38), an enzyme required for generation of cADPR, is found to be only associated with the inner membrane of the nuclear envelope (1). Although it is widely accepted that the nucleus also holds the apparatus required to produce nuclear InsP_3 , including PIP_2 as well as PLC (Ref. 65; reviewed in Ref. 27), and that PLC can be found associated with the nuclear membrane (28, 64), it is still not completely clear whether PIP_2 is localized in the nucleus. It has been proposed that nuclear PIP_2 could be confined within a lipid bilayer forming intranuclear membrane invaginations (51). Alternatively, there are reports that PIP_2 is located in the nucleoplasm rather than at the nuclear envelope (13, 78, 106). Nevertheless, there are several reports showing that nuclear PIP_2 hydrolysis involves the translocation of activated tyrosine kinase receptors from the plasma membrane to the nucleus (2, 20, 44, 89). For instance, IGF-1 (66), integrins (20), hepatocyte growth factor (44), and insulin (89) represent some agonists known to preferentially cause nuclear PIP_2 breakdown, producing InsP_3 and consequently Ca^{2+} signals in the nucleus. Moreover, it has been demonstrated that, upon growth factor stimulation, activation of nuclear PLC can occur through an additional pathway that involves relocation of MAP kinase to the nucleus (93). Even though the identification of several components of the Ca^{2+} toolkit within the nucleus represented solid evidence that could explain how Ca^{2+} is released from the nuclear envelope directly into the nucleoplasm, recognition and characterization of a structure named the nucleoplasmic reticulum (NR) (29) increased appreciation for the role of nuclear Ca^{2+} signaling, as it provided direct evidence of a compartment that stores Ca^{2+} deep in the nucleus and releases it within the nucleoplasm in a process that could occur completely independent of the cytosolic Ca^{2+} increases. The NR was shown to be continuous with the endoplasmic reticulum/nuclear envelope (29), as observed by lamin-A staining (FIGURE 1). Additional morphological characterization showed that the NR can be divided into two types (reviewed in Ref. 62). The type I NR is characterized by invagination of the inner nuclear membrane, whereas NR type II is formed by invagination of both the outer and inner nuclear membrane; and, although structurally different, both NR subtypes can coexist within a single nucleus (31, 32, 62). Indeed,

similar intranuclear extensions of the endoplasmic reticulum had been described previously and were thought to be widespread among mammalian cell types (32) and to be dynamic structures that become altered during cell proliferation or disease states (105). The NR has also been identified in plant cells, and it has been suggested that it regulates nuclear Ca^{2+} signaling in these cell types as well (22, 81). By showing that the NR gives rise to localized Ca^{2+} gradients in the nucleoplasm, a potential mechanism was revealed by which Ca^{2+} -dependent events can be regulated differentially in the nucleus, just as they are in the cytosol.

The NR was shown to express functional InsP_3R (29) and RyR (67), giving rise to local Ca^{2+} signals in the nuclear interior, triggered by either InsP_3 (29) or Ca^{2+} , respectively (67). To date, it is not known whether the TPCs or TRPC2 are also expressed along the NR. It is also not known whether and how the various intracellular Ca^{2+} channels in the NR interact to coordinate nuclear Ca^{2+} signals and




Figure 1. The nucleus contains the nucleoplasmic reticulum structure
 The nuclear envelope was labeled with an antibody against lamin-A (green) in SkHep1 (A), an adenocarcinoma liver cell line, and freshly isolated primary hepatocytes treated (B) or not treated (C) with staurosporine (STA) for 4 h, respectively. DNA was stained with propidium iodide. Note the presence of the nucleoplasmic reticulum (NR) in a cell line as well as in a primary culture. In the two top panels, the images represent three focal planes, depicting the three dimensionality of the NR. In the cell lineage used here, the NR seems to be constitutively present (A). In the primary culture of hepatocytes, the NR, not present under control condition, can be induced by drugs, such as STA (compare B with C), corroborating the findings that the NR is a dynamic and inducible structure. Bar in A = 5 μm ; bars in B and C = 10 μm .

cellular functions. Although more studies are needed to clarify these points, there are suggestions that the NR expresses InsP₃-kinase isoform B (IP3KB) (73), which can inactivate InsP₃ (26, 111) and SERCA (21), indicating that nuclear Ca²⁺ signals can be locally initiated and terminated within the nucleus.

Although the data described above demonstrate the potential independent behavior of the nucleus relative to the cytosol, they cannot rule out that, in response to some stimuli, the nuclear Ca²⁺ signaling machinery needs to be activated by intermediate effectors located in the cytosol. For instance, it was shown in a liver cell line that extracellular ATP

preferentially activates nuclear Ca²⁺ release via InsP₃ that diffuses from the cytosol to the nucleus to activate nuclear InsP₃R (59). Indeed, at present, different mechanisms for nuclear Ca²⁺ increases have been considered. One mechanism relies on the expression pattern of InsP₃R isoforms in the cell (FIGURE 2A). It involves cytosolic to nucleoplasmic InsP₃ diffusion to preferentially bind to the type II InsP₃R concentrated in the nucleus. The main question regarding this model is how InsP₃, as it diffuses across the cytosol, does not trigger Ca²⁺ release from the InsP₃R located in the endoplasmic reticulum. The answer may rely on the differing affinity of the various InsP₃R isoforms to InsP₃ as described above: InsP₃R-II > InsP₃R-I > InsP₃R-III (52, 74). Therefore, if type II InsP₃R expression is concentrated in the nucleus, Ca²⁺ signals are more likely to start there (59). A second mechanism consists of nuclear InsP₃ formation (FIGURE 2B). In support of this, it has been demonstrated that growth factors and their respective receptors translocate from the plasma membrane to the nucleus to stimulate the hydrolysis of nuclear PIP₂ with the formation of InsP₃ within the nucleus and consequently nuclear Ca²⁺ signals (2, 23, 44, 86, 89). Another possible mechanism is the stimulation of nuclear RyR (FIGURE 2C). This might occur when Ca²⁺ released from nuclear stores stimulates further Ca²⁺ release from RyR present along the NR. Although more experimental evidence is needed to support the importance of this pathway, the presence of RyR in the NR and its functionality was already described in C2C12 cells and in cardiomyocytes (45, 67). Finally, but no less important, is the model that predicts the juxtaposition of the components of the Ca²⁺ toolkit with the nuclear envelope. Recently, it was shown using cardiomyocytes that the invaginations of the plasma membrane, known as T-tubules, concentrate the machinery to generate InsP₃ in direct apposition to the components of the nuclear Ca²⁺ toolkit. Thus, after stimulation from an extracellular ligand, InsP₃ is generated in the vicinity of the nucleus diffusing into it and triggering independent nuclear Ca²⁺ signals (50).

These studies have clearly shown that the nucleus can be completely autonomous in terms of Ca²⁺ regulation. The presence of the Ca²⁺ signaling machinery in the nuclear interior has several implications, in physiological and pathological states, some of which will be described below.

Cellular Functions Mediated by Nuclear Ca²⁺

Ca²⁺ signaling regulates a variety of cellular functions, and, more recently, it has been shown that Ca²⁺ increases in the nucleoplasm regulate events




Figure 2. Working hypotheses to trigger nuclear Ca²⁺
 Different pathways to generate nuclear Ca²⁺ signals have been proposed. In 1 (left), the binding of an agonist to its transmembrane G-protein-coupled receptor (GPCR) produces InsP₃ (IP₃) in the cytosol. IP₃ diffuses to the nucleus and binds to the high-affinity, type II InsP₃R (IP₃R-II), located along the nucleoplasmic reticulum (NR), to release Ca²⁺ in the nucleoplasm. In 2 (middle), the binding of a growth factor to a receptor tyrosine kinase (TKR) may result in the translocation of the receptor to the nucleus to generate nuclear IP₃. IP₃ binds to the IP₃R to release Ca²⁺ in the nucleoplasm. The topology of internalized TRK is not known, and the cartoon shows TRK as being inside the vesicle merely for simplicity. In 3 (right), another putative pathway is initiated following the binding of a hormone to GPCR at the plasma membrane. The IP₃ generated within the cytosol can release Ca²⁺ from the endo-/sarcoplasmic reticulum, and Ca²⁺ may reach the nucleus to trigger Ca²⁺-induced Ca²⁺ release by activating the ryanodine receptor (RyR) in the NR.

that are distinct from the ones mediated by cytosolic Ca^{2+} (2, 6, 29, 43, 87, 89, 91). For instance, nuclear Ca^{2+} promotes translocation of protein kinase C (PKC) to the nuclear envelope, whereas cytosolic Ca^{2+} causes PKC translocation to the plasma membrane (29). Also, nuclear but not cytosolic Ca^{2+} generated by insulin receptor activation regulates hepatocyte proliferation after partial hepatectomy (2). This finding was in agreement with previous data that showed that buffering nuclear Ca^{2+} regulates the cell cycle by arresting cells in an early phase of mitosis (90). In fact, it was found that nuclear Ca^{2+} signals regulate the rate of cell proliferation both in vitro and in xenographic tumors (3, 89). One possible mechanism by which nuclear Ca^{2+} may control growth is by activating the promoter region of genes involved in cell proliferation (4). In addition to its direct effect on tumor growth, impaired nuclear Ca^{2+} signaling can sensitize adenocarcinoma cells to radiotherapy, in part by downregulating metalloproteinase and growth factor receptor expression and activation induced by X-ray irradiation (3).

Nuclear Ca^{2+} is also involved in cardiomyocyte hypertrophy. For example, buffering Ca^{2+} in the nucleus causes nuclear enlargement in neonatal cardiomyocytes, an early sign of cardiac hypertrophy (45). Nuclear Ca^{2+} activates the calcineurin/NFAT signaling cascade, culminating in increased ANP expression (45), a well known cardiomyocyte hypertrophic marker (88). However, cardiac hypertrophy is an adaptive response that can be induced in pathological and physiological conditions. Indeed, nuclear Ca^{2+} is indispensable for induction of hypertrophy in both health and disease. Regardless of the precise cellular compartment in which InsP_3 is produced, it must reach the nucleus interior to trigger nuclear Ca^{2+} -mediating hypertrophy (6).

Also dependent on the spatial properties of Ca^{2+} signals are transcriptional responses, since they are strongly influenced by the intracellular localization of Ca^{2+} transients. For example, neuronal gene expression is differentially controlled by nuclear and cytoplasmic Ca^{2+} signals: signaling pathways activated by cytoplasmic Ca^{2+} target the serum-response element (SRE), whereas increases in nuclear Ca^{2+} are critical for cyclic-AMP-response element (CRE)-dependent calcium-activated transcription (8, 47). Also, the transcriptional factor cyclic-AMP response element-binding protein (CREB) can function as a nuclear Ca^{2+} -responsive transcription factor (97) that is involved in mediating Ca^{2+} -dependent transcription of a number of genes, including c-fos (41). Nuclear Ca^{2+} buffering inhibits CREB-mediated gene transcription (47). In addition, nuclear Ca^{2+} controls, through nuclear calcium-/calmodulin (CaM)-dependent protein kinases, especially CaM Kinase IV, the activity of the coactivator CREB-binding protein

(CBP) (18). CBP interacts with CREB and with many transcription factors (42), a mechanism whereby nuclear Ca^{2+} can modulate the expression of different genes. Nuclear Ca^{2+} is similarly required for epidermal growth factor (EGF)-mediated transcriptional activation of Elk-1 (85). An additional target directly regulated by nuclear Ca^{2+} signaling is the downstream regulatory element antagonist modulator (DREAM) (69). All DREAM family members bind specifically to DNA and regulate transcription negatively. An increase in the concentration of nuclear Ca^{2+} causes DREAM to dissociate from its DNA binding sites and thereby allows the transcription of its target genes (58). On the other hand, nuclear Ca^{2+} also can negatively regulate the activity of transcription factors. For example, chelation of nuclear Ca^{2+} raises the activity of the transcription enhancer factor (TEAD) (103).

It is widely known that Ca^{2+} transients are crucial for synaptic activity and neuro-adaptations (reviewed in Ref. 7) including survival (114), memory consolidation (60), and pain (99). For example, histone deacetylases (HDACs) have a pivotal role linking external stimuli with gene expression. Inhibition of nuclear Ca^{2+} signaling interferes with the subcellular distribution of members of class IIa HDACs in hippocampal neurons, representing a novel transcriptional pathway regulated by nuclear Ca^{2+} (95). Also, synaptic activity is required for neuronal survival. Part of this neuroprotection is afforded by the resulting increase in Ca^{2+} in the cell nucleus due to activation of CREB (80) and inhibition of DREAM family members (58). Together, these changes increase the transcriptional activity of neurons, as already discussed in this review, and result in the triggering of a neuroprotective program referred to as Activity Inhibitor of Death (AID) (114). The core of this program consists of the robust induction of nuclear Ca^{2+} -related genes involved in survival and differentiation processes, including ATF3, GADD45 β , GADD45 γ , IFI202b, NPAS4, NR4A1, SERPINb2, Inhibin β (114), and BTG2 (113). Another important role of nuclear Ca^{2+} signaling in the nervous system is its relationship to pain. Inducing Ca^{2+} signals in the nucleoplasm increases the expression of genes that modulate neuronal excitation and morphology. In this case, neurons became more susceptible to painful stimuli, contributing to the amplification of nociceptive sensitivity (99).

Together, these data show that a large amount of information regarding nuclear Ca^{2+} signaling and cell function has been acquired in recent years. These findings not only contribute to our understanding of basic aspects of several physiological processes but also permit nuclear Ca^{2+} to be considered as a potential therapeutic target to treat diseases.

Future Perspectives for Nuclear Ca²⁺ Signaling

The characterization of the NR as a nuclear Ca²⁺ compartment, together with evidence demonstrating the presence of components of the components of the Ca²⁺ toolkit in the nuclear interior, add one more level of complexity to the already multifaceted field of Ca²⁺ signaling. A remaining intriguing question would be to understand whether and how the different intracellular Ca²⁺ channels present in the nucleus interact to decode Ca²⁺ signals to produce a specific cellular response. Moreover, the existence of micro-domains within the nucleus might also be important to tightly regulate nuclear Ca²⁺ signals and should be considered and investigated. ■

We thank Dr. Michael H. Nathanson (Yale University) for carefully reading the manuscript.

The authors acknowledge the supporting agencies CAPES, FAPEMIG, FAPEMIG-Pronex, CNPq, and HHMI. We also acknowledge the technical assistance of Gilson Nogueira.

No conflicts of interest, financial or otherwise, are declared by the author(s).

Author contributions: A.G.O. prepared figures; A.G.O., E.S.G., L.M.A., G.B.M., and M.F.L. drafted manuscript; A.G.O., G.B.M., and M.F.L. edited and revised manuscript; A.G.O., E.S.G., L.M.A., G.B.M., and M.F.L. approved final version of manuscript.

References

1. Adebajo OA, Anandatheerthavarada HK, Koval AP, Moonga BS, Biswas G, Sun L, Sodam BR, Bevis PJ, Huang CL, Epstein S, Lai FA, Avadhani NG, Zaidi M. A new function for CD38/ADP-ribosyl cyclase in nuclear Ca²⁺ homeostasis. *Nat Cell Biol* 1: 409–414, 1999.
2. Amaya MJ, Oliveira AG, Guimaraes ES, Casteluber MC, Carvalho SM, Andrade LM, Pinto MC, Mennone A, Oliveira CA, Resende RR, Menezes GB, Nathanson MH, Leite MF. The insulin receptor translocates to the nucleus to regulate cell proliferation in liver. *Hepatology* 59: 274–283, 2014.
3. Andrade LM, Geraldo JM, Gonçalves OX, Leite MTT, Catarina AM, Guimarães MM, Paes Leme AF, Yokoo S, Machado CR, Rajão MA, Carvalho SM, Gomes DA, Aguiar CJ, Souza-Fagundes EM, Zani CL, Resende RR, Martins-Filho OA, Leite MF. Nucleoplasmic calcium buffering sensitizes human squamous cell carcinoma to anticancer therapy. *J Cancer Sci Ther* 4: 131–139, 2012.
4. Andrade V, Guerra M, Jardim C, Melo F, Silva W, Ortega JM, Robert M, Nathanson MH, Leite F. Nucleoplasmic calcium regulates cell proliferation through legumain. *J Hepatol* 55: 626–635, 2011.
5. Anyatonwu GI, Ehrlich BE. Calcium signaling and polycystin-2. *Biochem Biophys Res Commun* 322: 1364–1373, 2004.
6. Arantes LA, Aguiar CJ, Amaya MJ, Figueiro NC, Andrade LM, Rocha-Resende C, Resende RR, Franchini KG, Guatimosim S, Leite MF. Nuclear inositol 1,4,5-trisphosphate is a necessary and conserved signal for the induction of both pathological and physiological cardiomyocyte hypertrophy. *J Mol Cell Cardiol* 53: 475–486, 2012.
7. Bading H. Nuclear calcium signalling in the regulation of brain function. *Nat Rev Neurosci* 14: 593–608, 2013.
8. Bading H, Hardingham GE, Johnson CM, Chawla S. Gene regulation by nuclear and cytoplasmic calcium signals. *Biochem Biophys Res Commun* 236: 541–543, 1997.
9. Berridge MJ. Calcium microdomains: organization and function. *Cell Calcium* 40: 405–412, 2006.

10. Berridge MJ, Lipp P, Bootman MD. The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* 1: 11–21, 2000.
11. Bers DM. Macromolecular complexes regulating cardiac ryanodine receptor function. *J Mol Cell Cardiol* 37: 417–429, 2004.
12. Blaustein MP, Lederer WJ. Sodium/calcium exchange: its physiological implications. *Physiol Rev* 79: 763–854, 1999.
13. Boronenkov IV, Loijens JC, Umeda M, Anderson RA. Phosphoinositide signaling pathways in nuclei are associated with nuclear speckles containing pre-mRNA processing factors. *Mol Biol Cell* 9: 3547–3560, 1998.
14. Brini M, Murgia M, Pasti L, Picard D, Pozzan T, Rizzuto R. Nuclear Ca²⁺ concentration measured with specifically targeted recombinant aequorin. *EMBO J* 12: 4813–4819, 1993.
15. Calcraft PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, Tang J, Rietdorf K, Teboul L, Chuang KT, Lin P, Xiao R, Wang C, Zhu Y, Lin Y, Wyatt CN, Parrington J, Ma J, Evans AM, Galione A, Zhu MX. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* 459: 596–600, 2009.
16. Cancela JM, Churchill GC, Galione A. Coordination of agonist-induced Ca²⁺-signalling patterns by NAADP in pancreatic acinar cells. *Nature* 398: 74–76, 1999.
17. Cardenas C, Escobar M, Garcia A, Osorio-Reich M, Hartel S, Foskett JK, Franzini-Armstrong C. Visualization of inositol 1,4,5-trisphosphate receptors on the nuclear envelope outer membrane by freeze-drying and rotary shadowing for electron microscopy. *J Struct Biol* 171: 372–381, 2010.
18. Chawla S, Hardingham GE, Quinn DR, Bading H. CBP: a signal-regulated transcriptional coactivator controlled by nuclear calcium and CaM kinase IV. *Science* 281: 1505–1509, 1998.
19. Choe CU, Ehrlich BE. The inositol 1,4,5-trisphosphate receptor (IP3R) and its regulators: sometimes good and sometimes bad teamwork. *Sci STKE* 2006: re15, 2006.
20. Clark EA, Brugge JS. Integrins and signal transduction pathways: the road taken. *Science* 268: 233–239, 1995.
21. Collado-Hilly M, Shirvani H, Jaillard D, Mauger JP. Differential redistribution of Ca²⁺-handling proteins during polarization of MDCK cells: effects on Ca²⁺ signalling. *Cell Calcium* 48: 215–224, 2010.
22. Collins DA, Carter CN, Rink JC, Scott AC, Wyatt CN, Allen NS. Plant nuclei can contain extensive grooves and invaginations. *Plant Cell* 12: 2425–2440, 2000.
23. De Angelis Campos AC, Rodrigues MA, de Andrade C, de Goes AM, Nathanson MH, Gomes DA. Epidermal growth factor receptors destined for the nucleus are internalized via a clathrin-dependent pathway. *Biochem Biophys Res Commun* 412: 341–346, 2011.
24. Decrock E, De Bock M, Wang N, Gadicherla AK, Bol M, Delvaeye T, Vandenaebelle P, Vinken M, Bultynck G, Krysko DV, Leybaert L. IP3, a small molecule with a powerful message. *Biochim Biophys Acta* 1833: 1772–1786, 2013.
25. Dellis O, Dedos SG, Tovey SC, Taufiq Ur R, Dubel SJ, Taylor CW. Ca²⁺ entry through plasma membrane IP3 receptors. *Science* 313: 229–233, 2006.
26. Dewaste V, Moreau C, De Smedt F, Bex F, De Smedt H, Wuytack F, Missiaen L, Erneux C. The three isoenzymes of human inositol-1,4,5-trisphosphate 3-kinase show specific intracellular localization but comparable Ca²⁺ responses on transfection in COS-7 cells. *Biochem J* 374: 41–49, 2003.
27. Divecha N, Banfic H, Irvine RF. Inositides and the nucleus and inositides in the nucleus. *Cell* 74: 405–407, 1993.
28. Divecha N, Banfic H, Irvine RF. The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase C to the nucleus. *EMBO J* 10: 3207–3214, 1991.
29. Echevarria W, Leite MF, Guerra MT, Zipfel WR, Nathanson MH. Regulation of calcium signals in the nucleus by a nucleoplasmic reticulum. *Nat Cell Biol* 5: 440–446, 2003.
30. Foskett JK, White C, Cheung KH, Mak DO. Inositol trisphosphate receptor Ca²⁺ release channels. *Physiol Rev* 87: 593–658, 2007.

31. Fricker M, Hollinshead M, White N, Vaux D. The convoluted nucleus. *Trends Cell Biol* 7: 181, 1997.
32. Fricker M, Hollinshead M, White N, Vaux D. Interphase nuclei of many mammalian cell types contain deep, dynamic, tubular membrane-bound invaginations of the nuclear envelope. *J Cell Biol* 136: 531–544, 1997.
33. Furuichi T, Furutama D, Hakamata Y, Nakai J, Takeshima H, Mikoshiba K. Multiple types of ryanodine receptor/ Ca^{2+} release channels are differentially expressed in rabbit brain. *J Neurosci* 14: 4794–4805, 1994.
34. Galione A, Morgan AJ, Arredouani A, Davis LC, Rietdorf K, Ruas M, Parrington J. NAADP as an intracellular messenger regulating lysosomal calcium-release channels. *Biochem Soc Trans* 38: 1424–1431, 2010.
35. Gerasimenko JV, Lur G, Sherwood MW, Ebisui E, Tepikin AV, Mikoshiba K, Gerasimenko OV, Petersen OH. Pancreatic protease activation by alcohol metabolite depends on Ca^{2+} release via acid store IP₃ receptors. *Proc Natl Acad Sci USA* 106: 10758–10763, 2009.
36. Gerasimenko JV, Maruyama Y, Yano K, Dolman NJ, Tepikin AV, Petersen OH, Gerasimenko OV. NAADP mobilizes Ca^{2+} from a thapsigargin-sensitive store in the nuclear envelope by activating ryanodine receptors. *J Cell Biol* 163: 271–282, 2003.
37. Gerasimenko O, Gerasimenko J. New aspects of nuclear calcium signalling. *J Cell Sci* 117: 3087–3094, 2004.
38. Gerasimenko OV, Gerasimenko JV, Petersen OH, Tepikin AV. Short pulses of acetylcholine stimulation induce cytosolic Ca^{2+} signals that are excluded from the nuclear region in pancreatic acinar cells. *Pflügers Arch* 432: 1055–1061, 1996.
39. Gerasimenko OV, Gerasimenko JV, Tepikin AV, Petersen OH. ATP-dependent accumulation and inositol trisphosphate- or cyclic ADP-ribose-mediated release of Ca^{2+} from the nuclear envelope. *Cell* 80: 439–444, 1995.
40. Giannini G, Conti A, Mammarella S, Scrobogna M, Sorrentino V. The ryanodine receptor/calcium channel genes are widely and differentially expressed in murine brain and peripheral tissues. *J Cell Biol* 128: 893–904, 1995.
41. Ginty DD, Bonni A, Greenberg ME. Nerve growth factor activates a Ras-dependent protein kinase that stimulates c-fos transcription via phosphorylation of CREB. *Cell* 77: 713–725, 1994.
42. Goldman PS, Tran VK, Goodman RH. The multifunctional role of the co-activator CBP in transcriptional regulation. *Recent Prog Horm Res* 52: 103–119; discussion 119–120, 1997.
43. Gomes DA, Leite MF, Bennett AM, Nathanson MH. Calcium signaling in the nucleus. *Can J Physiol Pharmacol* 84: 325–332, 2006.
44. Gomes DA, Rodrigues MA, Leite MF, Gomez MV, Varnai P, Balla T, Bennett AM, Nathanson MH. c-Met must translocate to the nucleus to initiate calcium signals. *J Biol Chem* 283: 4344–4351, 2008.
45. Guatimosim S, Amaya MJ, Guerra MT, Aguiar CJ, Goes AM, Gomez-Viquez NL, Rodrigues MA, Gomes DA, Martins-Cruz J, Lederer WJ, Leite MF. Nuclear Ca^{2+} regulates cardiomyocyte function. *Cell Calcium* 44: 230–242, 2008.
46. Hakamata Y, Nakai J, Takeshima H, Imoto K. Primary structure and distribution of a novel ryanodine receptor/calcium release channel from rabbit brain. *FEBS Lett* 312: 229–235, 1992.
47. Hardingham GE, Chawla S, Johnson CM, Bading H. Distinct functions of nuclear and cytoplasmic calcium in the control of gene expression. *Nature* 385: 260–265, 1997.
48. Hennager DJ, Welsh MJ, DeLisle S. Changes in either cytosolic or nucleoplasmic inositol 1,4,5-trisphosphate levels can control nuclear Ca^{2+} concentration. *J Biol Chem* 270: 4959–4962, 1995.
49. Humbert JP, Matter N, Artault JC, Koppler P, Malviya AN. Inositol 1,4,5-trisphosphate receptor is located to the inner nuclear membrane vindicating regulation of nuclear calcium signaling by inositol 1,4,5-trisphosphate. Discrete distribution of inositol phosphate receptors to inner and outer nuclear membranes. *J Biol Chem* 271: 478–485, 1996.
50. Ibarra C, Vicencio JM, Estrada M, Lin Y, Rocco P, Rebellato P, Munoz JP, Garcia-Prieto J, Quest AF, Chiong M, Davidson SM, Bulatovic I, Grinemo KH, Larsson O, Szabadkai G, Uhlen P, Jaimovich E, Lavandero S. Local control of nuclear calcium signaling in cardiac myocytes by perinuclear microdomains of sarcolemmal insulin-like growth factor 1 receptors. *Circ Res* 112: 236–245, 2013.
51. Irvine RF. Nuclear inositide signalling: expansion, structures and clarification. *Biochim Biophys Acta* 1761: 505–508, 2006.
52. Iwai M, Michikawa T, Bosanac I, Ikura M, Mikoshiba K. Molecular basis of the isoform-specific ligand-binding affinity of inositol 1,4,5-trisphosphate receptors. *J Biol Chem* 282: 12755–12764, 2007.
53. Kottgen M, Walz G. Subcellular localization and trafficking of polycystins. *Pflügers Arch* 451: 286–293, 2005.
54. Koulen P, Cai Y, Geng L, Maeda Y, Nishimura S, Witzgall R, Ehrlich BE, Somlo S. Polycystin-2 is an intracellular calcium release channel. *Nat Cell Biol* 4: 191–197, 2002.
55. Lai FA, Dent M, Wickenden C, Xu L, Kumari G, Misra M, Lee HB, Sar M, Meissner G. Expression of a cardiac Ca^{2+} -release channel isoform in mammalian brain. *Biochem J* 288: 553–564, 1992.
56. Lanini L, Bachs O, Carafoli E. The calcium pump of the liver nuclear membrane is identical to that of endoplasmic reticulum. *J Biol Chem* 267: 11548–11552, 1992.
57. Lanner JT, Georgiou DK, Joshi AD, Hamilton SL. Ryanodine receptors: structure, expression, molecular details, and function in calcium release. *Cold Spring Harb Perspect Biol* 2: a003996, 2010.
58. Ledo F, Kremer L, Mellstrom B, Naranjo JR. Ca^{2+} -dependent block of CREB-CBP transcription by repressor DREAM. *EMBO J* 21: 4583–4592, 2002.
59. Leite MF, Thrower EC, Echevarria W, Koulen P, Hirata K, Bennett AM, Ehrlich BE, Nathanson MH. Nuclear and cytosolic calcium are regulated independently. *Proc Natl Acad Sci USA* 100: 2975–2980, 2003.
60. Limback-Stokin K, Korzus E, Nagaoka-Yasuda R, Mayford M. Nuclear calcium/calmodulin regulates memory consolidation. *J Neurosci* 24: 10858–10867, 2004.
61. Lipp P, Thomas D, Berridge MJ, Bootman MD. Nuclear calcium signalling by individual cytoplasmic calcium puffs. *EMBO J* 16: 7166–7173, 1997.
62. Malhas A, Goulbourne C, Vaux DJ. The nucleoplasmic reticulum: form and function. *Trends Cell Biol* 21: 362–373, 2011.
63. Malviya AN, Rogue P, Vincendon G. Stereospecific inositol 1,4,5-³²P]trisphosphate binding to isolated rat liver nuclei: evidence for inositol trisphosphate receptor-mediated calcium release from the nucleus. *Proc Natl Acad Sci USA* 87: 9270–9274, 1990.
64. Manzoli L, Martelli AM, Billi AM, Faenza I, Fiume R, Cocco L. Nuclear phospholipase C: involvement in signal transduction. *Prog Lipid Res* 44: 185–206, 2005.
65. Maraldi NM, Cocco L, Capitani S, Mazzotti G, Barnabei O, Manzoli FA. Lipid-dependent nuclear signalling: morphological and functional features. *Adv Enzyme Regul* 34: 129–143, 1994.
66. Maraldi NM, Zini N, Ognibene A, Martelli AM, Barbieri M, Mazzotti G, Manzoli FA. Immunocytochemical detection of the intranuclear variations of phosphatidylinositol 4,5-bisphosphate amount associated with changes of activity and amount of phospholipase C beta 1 in cells exposed to mitogenic or differentiating agonists. *Biol Cell* 83: 201–210, 1995.
67. Marius P, Guerra MT, Nathanson MH, Ehrlich BE, Leite MF. Calcium release from ryanodine receptors in the nucleoplasmic reticulum. *Cell Calcium* 39: 65–73, 2006.
68. McPherson PS, Campbell KP. Characterization of the major brain form of the ryanodine receptor/ Ca^{2+} release channel. *J Biol Chem* 268: 19785–19790, 1993.
69. Mellstrom B, Savignac M, Gomez-Villafuertes R, Naranjo JR. Ca^{2+} -operated transcriptional networks: molecular mechanisms and in vivo models. *Physiol Rev* 88: 421–449, 2008.
70. Meszaros LG, Bak JZ, Chu A. Cyclic ADP-ribose as an endogenous regulator of the non-skeletal type ryanodine receptor Ca^{2+} channel. *Nature* 364: 76–79, 1993.
71. Nakai J, Imagawa T, Hakamat Y, Shigekawa M, Takeshima H, Numa S. Primary structure and functional expression from cDNA of the cardiac ryanodine receptor/calcium release channel. *FEBS Lett* 271: 169–177, 1990.
72. Nakanishi S, Kuwajima G, Mikoshiba K. Immunohistochemical localization of ryanodine receptors in mouse central nervous system. *Neurosci Res* 15: 130–142, 1992.
73. Nalaskowski MM, Fliegert R, Ernst O, Brehm MA, Fanick W, Windhorst S, Lin H, Giehler S, Hein J, Lin YN, Mayr GW. Human inositol 1,4,5-trisphosphate 3-kinase isoform B (IP3KB) is a nucleocytoplasmic shuttling protein specifically enriched at cortical actin filaments and at invaginations of the nuclear envelope. *J Biol Chem* 286: 4500–4510, 2011.
74. Newton CL, Mignery GA, Sudhof TC. Co-expression in vertebrate tissues and cell lines of multiple inositol 1,4,5-trisphosphate (InsP₃) receptors with distinct affinities for InsP₃. *J Biol Chem* 269: 28613–28619, 1994.
75. Neylon CB, Richards SM, Larsen MA, Agrotis A, Bobik A. Multiple types of ryanodine receptor/ Ca^{2+} release channels are expressed in vascular smooth muscle. *Biochem Biophys Res Commun* 215: 814–821, 1995.
76. Nicotera P, McConkey DJ, Jones DP, Orrenius S. ATP stimulates Ca^{2+} uptake and increases the free Ca^{2+} concentration in isolated rat liver nuclei. *Proc Natl Acad Sci USA* 86: 453–457, 1989.
77. Nicotera P, Orrenius S, Nilsson T, Berggren PO. An inositol 1,4,5-trisphosphate-sensitive Ca^{2+} pool in liver nuclei. *Proc Natl Acad Sci USA* 87: 6858–6862, 1990.
78. Osborne SL, Thomas CL, Gschmeissner S, Schiavo G. Nuclear PtdIns(4,5)P₂ assembles in a mitotically regulated particle involved in pre-mRNA splicing. *J Cell Sci* 114: 2501–2511, 2001.
79. Otsu K, Willard HF, Khanna VK, Zorzato F, Green NM, MacLennan DH. Molecular cloning of cDNA encoding the Ca^{2+} release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic reticulum. *J Biol Chem* 265: 13472–13483, 1990.
80. Papadia S, Stevenson P, Hardingham NR, Bading H, Hardingham GE. Nuclear Ca^{2+} and the cAMP response element-binding protein family mediate a late phase of activity-dependent neuroprotection. *J Neurosci* 25: 4279–4287, 2005.

81. Pauly N, Knight MR, Thuleau P, van der Luit AH, Moreau M, Trewavas AJ, Ranjeva R, Mazars C. Control of free calcium in plant cell nuclei. *Nature* 405: 754–755, 2000.
82. Perocchi F, Gohil VM, Girgis HS, Bao XR, McCombs JE, Palmer AE, Mootha VK. MICU1 encodes a mitochondrial EF hand protein required for Ca^{2+} uptake. *Nature* 467: 291–296, 2010.
83. Pinton P, Pozzan T, Rizzuto R. The Golgi apparatus is an inositol 1,4,5-trisphosphate-sensitive Ca^{2+} store, with functional properties distinct from those of the endoplasmic reticulum. *EMBO J* 17: 5298–5308, 1998.
84. Pozzan T, Rizzuto R, Volpe P, Meldolesi J. Molecular and cellular physiology of intracellular calcium stores. *Physiol Rev* 74: 595–636, 1994.
85. Pust T, Wu JJ, Zimmerman TL, Zhang L, Ehrlich BE, Berchtold MW, Hoek JB, Karpen SJ, Nathanson MH, Bennett AM. Epidermal growth factor-mediated activation of the ETS domain transcription factor Elk-1 requires nuclear calcium. *J Biol Chem* 277: 27517–27527, 2002.
86. Reilly JF, Maher PA. Importin beta-mediated nuclear import of fibroblast growth factor receptor: role in cell proliferation. *J Cell Biol* 152: 1307–1312, 2001.
87. Resende RR, Andrade LM, Oliveira AG, Guimaraes ES, Guatimosim S, Leite MF. Nucleoplasmic calcium signaling and cell proliferation: calcium signaling in the nucleus. *Cell Commun Signal* 11: 14, 2013.
88. Roderick HL, Higazi DR, Smyrniats I, Fearnley C, Harzheim D, Bootman MD. Calcium in the heart: when it's good, it's very very good, but when it's bad, it's horrid. *Biochem Soc Trans* 35: 957–961, 2007.
89. Rodrigues MA, Gomes DA, Andrade VA, Leite MF, Nathanson MH. Insulin induces calcium signals in the nucleus of rat hepatocytes. *Hepatology* 48: 1621–1631, 2008.
90. Rodrigues MA, Gomes DA, Leite MF, Grant W, Zhang L, Lam W, Cheng YC, Bennett AM, Nathanson MH. Nucleoplasmic calcium is required for cell proliferation. *J Biol Chem* 282: 17061–17068, 2007.
91. Rodrigues MA, Gomes DA, Nathanson MH, Leite MF. Nuclear calcium signaling: a cell within a cell. *Braz J Med Biol Res* 42: 17–20, 2009.
92. Ruas M, Rietdorf K, Arredouani A, Davis LC, Lloyd-Evans E, Koegel H, Funnell TM, Morgan AJ, Ward JA, Watanabe K, Cheng X, Churchill GC, Zhu MX, Platt FM, Wessel GM, Parrington J, Galione A. Purified TPC isoforms form NAADP receptors with distinct roles for Ca^{2+} signaling and endolysosomal trafficking. *Curr Biol* 20: 703–709, 2010.
93. Santella L, Carafoli E. Calcium signaling in the cell nucleus. *FASEB J* 11: 1091–1109, 1997.
94. Santella L, Kyozuka K. Effects of 1-methyladenine on nuclear Ca^{2+} transients and meiosis resumption in starfish oocytes are mimicked by the nuclear injection of inositol 1,4,5-trisphosphate and cADP-ribose. *Cell Calcium* 22: 11–20, 1997.
95. Schlumm F, Mauceri D, Freitag HE, Bading H. Nuclear calcium signaling regulates nuclear export of a subset of class IIa histone deacetylases following synaptic activity. *J Biol Chem* 288: 8074–8084, 2013.
96. Sharp AH, McPherson PS, Dawson TM, Aoki C, Campbell KP, Snyder SH. Differential immunohistochemical localization of inositol 1,4,5-trisphosphate- and ryanodine-sensitive Ca^{2+} release channels in rat brain. *J Neurosci* 13: 3051–3063, 1993.
97. Sheng M, Thompson MA, Greenberg ME. CREB: a Ca^{2+} -regulated transcription factor phosphorylated by calmodulin-dependent kinases. *Science* 252: 1427–1430, 1991.
98. Shirakawa H, Miyazaki S. Spatiotemporal analysis of calcium dynamics in the nucleus of hamster oocytes. *J Physiol* 494: 29–40, 1996.
99. Simonetti M, Hagenston AM, Vardeh D, Freitag HE, Mauceri D, Lu J, Satagopam VP, Schneider R, Costigan M, Bading H, Kuner R. Nuclear calcium signaling in spinal neurons drives a genomic program required for persistent inflammatory pain. *Neuron* 77: 43–57, 2013.
100. Sonleitner A, Conti A, Bertocchini F, Schindler H, Sorrentino V. Functional properties of the ryanodine receptor type 3 (RyR3) Ca^{2+} release channel. *EMBO J* 17: 2790–2798, 1998.
101. Stehno-Bittel L, Luckhoff A, Clapham DE. Calcium release from the nucleus by InsP3 receptor channels. *Neuron* 14: 163–167, 1995.
102. Taylor CW, Dale P. Intracellular Ca^{2+} channels: a growing community. *Mol Cell Endocrinol* 353: 21–28, 2012.
103. Thompson M, Andrade VA, Andrade SJ, Pust T, Ortega JM, Goes AM, Leite MF. Inhibition of the TEF/TEAD transcription factor activity by nuclear calcium and distinct kinase pathways. *Biochem Biophys Res Commun* 301: 267–274, 2003.
104. Tsiokas L, Kim S, Ong EC. Cell biology of polycystin-2. *Cell Signal* 19: 444–453, 2007.
105. Wamhoff BR, Dixon JL, Sturek M. Atorvastatin treatment prevents alterations in coronary smooth muscle nuclear Ca^{2+} signaling in diabetic dyslipidemia. *J Vasc Res* 39: 208–220, 2002.
106. Watt SA, Kular G, Fleming IN, Downes CP, Lucocq JM. Subcellular localization of phosphatidylinositol 4,5-bisphosphate using the pleckstrin homology domain of phospholipase C delta1. *Biochem J* 363: 657–666, 2002.
107. Waybill MM, Yelamarty RV, Zhang Y, Scaduto RC, LaNoue KF, Hsu CJ, Smith BC, Tillotson DL, Yu FTS, Cheung JY. Nuclear calcium gradients in cultured rat hepatocytes. *Am J Physiol Endocrinol Metab* 261: E49–E57, 1991.
108. Wojcikiewicz RJ. Type I, II, and III inositol 1,4,5-trisphosphate receptors are unequally susceptible to down-regulation and are expressed in markedly different proportions in different cell types. *J Biol Chem* 270: 11678–11683, 1995.
109. Wu G, Xie X, Lu ZH, Ledeen RW. Sodium-calcium exchanger complexed with GM1 ganglioside in nuclear membrane transfers calcium from nucleoplasm to endoplasmic reticulum. *Proc Natl Acad Sci USA* 106: 10829–10834, 2009.
110. Xie X, Wu G, Lu ZH, Ledeen RW. Potentiation of a sodium-calcium exchanger in the nuclear envelope by nuclear GM1 ganglioside. *J Neurochem* 81: 1185–1195, 2002.
111. Yu JC, Lloyd-Burton SM, Irvine RF, Schell MJ. Regulation of the localization and activity of inositol 1,4,5-trisphosphate 3-kinase B in intact cells by proteolysis. *Biochem J* 392: 435–441, 2005.
112. Zalk R, Lehnart SE, Marks AR. Modulation of the ryanodine receptor and intracellular calcium. *Annu Rev Biochem* 76: 367–385, 2007.
113. Zhang SJ, Steijaert MN, Lau D, Schutz G, Delucinge-Vivier C, Descombes P, Bading H. Decoding NMDA receptor signaling: identification of genomic programs specifying neuronal survival and death. *Neuron* 53: 549–562, 2007.
114. Zhang SJ, Zou M, Lu L, Lau D, Ditzel DA, Delucinge-Vivier C, Aso Y, Descombes P, Bading H. Nuclear calcium signaling controls expression of a large gene pool: identification of a gene program for acquired neuroprotection induced by synaptic activity. *PLoS Genet* 5: e1000604, 2009.